

**REGISTRY OF PATENTS
SINGAPORE**


This is to certify that the annexed is a true copy of specification as filed for the following Singapore patent application.

Date of Filing : 24 FEBRUARY 2001

Application Number : 200100727-7

Applicant(s) /
Proprietor(s) of
Patent : URAH RESOURCES (NIGERIA) LTD;
OLOBO JONATHAN OBAJE

Title of Invention : SOLVENT-FREE TRANS-ACIDOLYSIS
PROCESS FOR THE PREPARATION OF
EDIBLE SURFACE-ACTIVE
CARBOHYDRATE FATTY-ACID ESTERS



SHARMAINE WU (Ms)

Assistant Registrar
for REGISTRAR OF PATENTS



24 FEB 2001
200100727-7

Rule 19


SINGAPORE
PATENTS ACT
(CHAPTER 221)
PATENTS RULESThe Registrar of Patents
Registry of PatentsREQUEST FOR THE GRANT OF A PATENT
THE GRANT OF A PATENT IS REQUESTED BY THE UNDERSIGNED ON THE BASIS OF THE PRESENT APPLICATION

I. Title of Invention	SOLVENT-FREE TRANS-ACIDOLYSIS PROCESS FOR THE PREPARATION OF EDIBLE SURFACE- ACTIVE CARBOHYDRATE FATTY-ACID ESTERS	
II. Applicant(s) (See note 2)	(a) Name	Olobo Jonathan Obaje Tel: (65) 7744482
	Body Description/ Residency	1. Singapore (ordinary Residence with EP) 2. Nigeria: Citizen
	Street Name & Number	Block 510 #04-311, West Coast Drive
	City	Singapore 120510
	State	Singapore
	Country	Singapore
	(b) Name	N.A.
	Body Description/ Residency	
	Street Name & Number	
	City	
	State	
	Country	
	(c) Name	N.A.
	Body Description/ Residency	
	State	
	Country	

#160

24/2

R 14339



III. Declaration of Priority (see note 3)	Country/Country Designated		File no.		
	Filing Date				
	Country/Country Designated		File no.		
	Filing Date				
IV. Inventors (See note 4)					
(a) The applicant(s) is/are the sole/joint inventor(s).		<div style="display: flex; justify-content: space-around;"> <div> <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No </div> <div> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No </div> </div>			
(b) A statement on Patents Form 8 is/will be furnished.					
V. Name of Agent (if any) (See note 5)					
VI. Address for Service (See note 6)		Block No	510	Level No	04
		Unit No	311	Postal Code	120510
		Street Name	West	Coast	Drive
		Building Name	Blk 510		
VII. Claiming an earlier filing date under section 20(3), 26(6) or 47(4). (See note 7)		Application No			
		Filing Date			
		[Please tick in the relevant space provided]: () Proceeding under rule 27(1)(a). Date on which the earlier application was amended = _____ or () Proceeding under rule 27(1)(b).			

200100727-7

24 FEB 2001

VIII. Invention has been displayed at an International Exhibition (See note 8)		<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	
IX. Section 114 requirements (See note 9)		The invention relates to and/or used a micro-organism deposited for the purposes of disclosure in accordance with section 114 with a depository authority under the Budapest Treaty. <input type="checkbox"/> Yes <input type="checkbox"/> No	
X. Check List (To be filled in by applicant or agent)	A. The application contains the following number of sheet(s):-		
	1. Request	4	sheets
	2. Description	11	sheets
	3. Claim(s).	2	sheets
	4. Drawing(s).	—	sheets
	5. Abstract.	1	sheets
	B. The application as filed is accompanied by:-		
	1. Priority document	<input type="checkbox"/>	
	2. Translation of priority document	<input type="checkbox"/>	
	3. Statement of Inventorship & right to grant	<input type="checkbox"/>	
4. International Exhibition Certificate	<input type="checkbox"/>		
XI. Signature(s) (See note 10)	Applicant (a)	OLDBO JONATHAN OBAJE <i>Johny</i>	
	Date	24 Feb 2001	
	Applicant (b)		
	Date		
	Applicant (c)		
	Date		

ENCLOSED US DISCLOSURE DOCUMENT NO 475021
Dated JUNE 01 2000.

NOTES:

200100727-7

24 FEB 2001

1. This form when completed, should be brought or sent to the Registry of Patents together with the prescribed fee and 3 copies of the description of the invention, and of any drawings.
2. Enter the name and address of each applicant in the spaces provided at paragraph II. Names of individuals should be indicated in full and the surname or family name should be underlined. The names of all partners in a firm must be given in full. The place of residence of each individual should also be furnished in the space provided. Bodies corporate should be designated by their corporate name and country of incorporation and, where appropriate, the state of incorporation within that country should be entered where provided. Where more than 3 applicants are to be named, the names and address of the fourth and any further applicants should be given on a separate sheet attached to this form together with the signature of each of these further applicants.
3. The declaration of priority at paragraph III should state the date of the previous filing, the country in which it was made, and indicate the file number, if available. Where the application relied upon in an International Application or a regional patent application e.g. European patent application, one of the countries designated in that application [being one falling under the Patents (Convention Countries) Order] should be identified and the name of that country should be entered in the space provided.
4. Where the applicant or applicants is/are the sole inventor or the joint inventors, paragraph IV should be completed by marking the 'YES' Box in the declaration (a) and the 'NO' Box in the alternative statement (b). Where this is not the case, the 'NO' Box in declaration (a) should be marked and a statement will be required to be filed on Patents Form 8.
5. If the applicant has appointed an agent to act on his behalf, the agent's name should be indicated in the spaces available at paragraph V.
6. An address for service in Singapore to which all documents may be sent must be stated at paragraph VI. It is recommended that a telephone number be provided if an agent is not appointed.
7. When an application is made by virtue of section 20(3), 26(6) or 47(4), the appropriate section should be identified at paragraph VII and the number of the earlier application or any patent granted thereon identified. Applicants proceeding under section 26(6) should identify which provision in rule 27 they are proceeding under. If the applicants are proceeding under rule 27(1)(a), they should also indicate the date on which the earlier application was amended.
8. Where the applicant wishes an earlier disclosure of the invention by him at an International Exhibition to be disregarded in accordance with section 14(4)(c), then the 'YES' Box at paragraph VIII should be marked. Otherwise the 'NO' Box should be marked.
9. Where in disclosing the invention the application refers to one or more micro-organisms deposited with a depository authority under the Budapest Treaty, then the 'YES' Box at paragraph IX should be marked. Otherwise the 'NO' Box should be marked.
10. Attention is drawn to rules 90 and 105 of the Patent Rules. Where there are more than 3 applicants, see also Note 2 above.
11. Applicants resident in Singapore are reminded that if the Registry of Patents considers that an application contains information the publication of which might be prejudicial to the defence of Singapore or the safety of the public, it may prohibit or restrict its publication or communication. Any person resident in Singapore and wishing to apply for patent protection in other countries must first obtain permission from the Singapore Registry of Patents unless they have already applied for a patent for the same invention in Singapore. In the latter case, no application should be made overseas until at least 2 months after the application has been filed in Singapore.

For Official Use

Application Filing Date: / /

Request received on : / /

Fee received on : / /

Amount :

*Cash/Cheque/Money Order No:

*Delete whichever is inapplicable

TITLE OF THE INVENTION
SOLVENT-FREE TRANS-ACIDOLYSIS PROCESS FOR THE
PREPARATION OF EDIBLE SURFACE-ACTIVE CARBOHYDRATE
FATTY-ACID ESTERS

5

BACKGROUND OF THE INVENTION

(a) Field of the Invention

The present invention relates to process for production of edible
10 carbohydrate fatty-acid esters, and in particular, a low temperature,
solvent-free trans-acidolysis reaction between carbohydrate ester of
low molecular-weight carboxylic acid and free fatty-acid under
reduced pressure and in the presence of an acid catalyst.

15 (b) Description of the Prior Art

Carbohydrate fatty-acid esters are odorless, non-irritant nonionic
surfactants with excellent biodegradability properties. They are used
as low calorie fat substitute, or to solubilize membrane proteins, and
20 to formulate many grades of detergents, pharmaceutical, food and
cosmetic products. Carbohydrate fatty-acid esters also exhibit anti-
microbial and insecticidal activities and are used as cerebral
metabolism improving agent, and in the treatment of colonic
disorders. But the synthetic methods for carbohydrates fatty-acid
25 esters are faced with many limitations.

U.S. Patent No. 5,945,519 describes a process for the solvent-free
preparation of sucrose fatty acid esters and their mixtures with

This Sheet Has Been Replaced



nonsugar polyol fatty acid esters. In this process, a sucrose is reacted with one or more fatty acid alkyl esters of a chain length of 6 to 20 carbon atoms and the reaction mixture is then reacted at reduced pressure with fatty acid alkyl ester, and then filtered without
5 addition of solvent.

U.S. Patent No. 4,614,718 discloses the synthesis of sugar or sugar-alcohol fatty acid esters by incubating an aqueous mixture of a sugar or sugar-alcohol, a C8 – C22 fatty acid and a catalytically
10 active amount of a lipolytic enzyme derived from the genus Pseudomonas, Enterobacterium, Chromobacterium, Mucor or Candida at a temperature from about 20 degree C to about 60 degree C, and recovering the resulting ester from the mixture.

15 U.S. Patent No. 5,908,922 (issued to Kasseri, et al, June 1, 1999) discloses a method for producing a sucrose fatty acid ester, which comprises subjecting sucrose and a lower alkyl ester of fatty acid to an ester exchanger reaction in an organic solvent in the presence of an alkali catalyst to produce the corresponding sucrose fatty acid
20 ester.

U.S. Patent No. 4,927,920 (issued to Wagner, et al, 22 May 1990) discloses a method for making sugar esters by reacting a sugar and an ester of the sugar having a degree of substitution greater than
25 about 2 in an organic solvent in the presence of a transesterification catalyst to form a sugar ester having a degree of substitution up to 2;

This Sheet Has Been Replaced



and recovering the sugar ester having a degree of substitution up to 2 at a temperature below the boiling point of the organic solvent.

U.S. Patent No. 4,996,309 discloses a process for preparing
5 sucrose fatty acid esters by reacting sucrose and a fatty acid alkyl ester in an aqueous reaction system in the presence of a catalyst, adding water to the reaction mixture to dissolve it, adjusting the reaction mixture to a neutral pH region, adding a neutral salt to the solution to precipitate the sucrose fatty acid esters, separating and
10 washing the precipitate with an acidic water, and subjecting the washing liquid to ultrafiltration.

The conventional methods for producing carbohydrate fatty esters are encountered with the following disadvantages and limitations:

- 15 (1) the transesterification method, currently favoured by most industrial producers of carbohydrate fatty esters, is a reversible process with poor conversion equilibria because the substrate sugars and fatty acid moieties are immiscible due to polarity differences. Thus, low product yield
20 characterize the processes;
- (2) Mutual solvents, such as pyridine, N, N-dimethylformamide (DMF), dimethylacetamide (DMA), dimethylsulfoxide (DMSO), chloroform, benzene and toluene, are required to solubilize the substrates. These mutual solvents are toxic
25 and cannot be removed to the level compatible with current regulations, thus limiting product applications.



- (3) High process cost resulting from high-energy cost and complex purification procedures.
- (4) Thermal degradation and isomerization of the reactant carbohydrates at temperatures above 100 degree C, which is normally required in most known processes.

It is a principal object of the present invention to provide a solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters, which obviates the above drawbacks and/or limitations.

SUMMARY OF THE INVENTION

Accordingly, it is an object of the present invention to provide a solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters, wherein the process comprises a low temperature, solvent-free trans-acidolysis reaction between carbohydrate ester of low molecular-weight carboxylic acid (C2 to C3) and free fatty acid (C6 to C22) under reduced pressure and in the presence of an acid catalyst.

Yet another object of the present invention is to provide a solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters, wherein mono-, di- and poly-fatty esters of C2 and C3 acylated carbohydrates, with hydrophile-lipophile balance (HLB) values of 2 to 10, are obtained depending on the initial molar ratio of the reactants and chain-length of the fatty acid moiety.

This Sheet Has Been Replaced



A further object of the present invention is to provide a solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters, wherein acid or base catalyzed partial deacylation lead to carbohydrate fatty esters with free hydroxyl groups, and having HLB values of 8-18.

Yet a further object of the present invention is to provide a solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters, wherein the unreacted free fatty acids and carbohydrate esters of low molecular-weight carboxylic acids, which are removed during purification and recycled to the starting reactant mixture.

The advantages of the process in accordance with the present invention, which circumvent the above limitations, are

- (1) avoiding solvent as a reaction medium. The C2-C3 acyl group attached to the reactant carbohydrate is a good protecting and leaving group and, at the same time, enhances the solubility of the carbohydrate moiety in the fatty acid;
- (2) using low pressure to drive the reaction equilibrium forward at a reduced energy demand. This results in high yield (>90), less isomerization and degradation of products.
- (3) The feedstock is renewable natural resources, readily available and cheap. Unreacted substrates are recycled. This process is thus commercially viable.

This Sheet Has Been Replaced



- (4) The invention is applicable to both batch-wise carbohydrate fatty ester synthesis and to continuous synthesis.
- (5) High-grade (98% pure) carbohydrate fatty acid esters can be obtained.
- (6) This process is adaptable for producing a number carbohydrate fatty acid esters.

Other objects and advantages of the invention will become apparent upon reading the following detailed description.

DETAILED DESCRIPTION OF THE PRESENT INVENTION


The present invention will be described in detail.

In accordance with the solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters of the present invention, acylated carbohydrate with chain-length C2-C3 includes mono to trisaccharide carbohydrates [see molecular structure], and the free fatty acid has chain-length C6 – C22.

In the present process, the acid catalyst includes concentrated sulphuric acid and alky sulfonic acid polysiloxanes. The carbohydrate fatty-acid esters of the present process has the following molecular structures:

This Sheet Has Been Replaced






wherein R2, R3, R5 are each selected from the group consisting of H, CH₃CO, CH₃CH₂CO, and C6-C22 fatty acyl group; R1, and R4 are each selected from the group consisting of a partially or peracylated mono- or di-saccharide, H, CH₃CO, CH₃CH₂CO or C6-C22 fatty acyl group; and n is an integer with value equal to 1, 2, or 3; wherein the primary monosaccharide unit, i.e. the [---]_n unit, could be a furanosyl, pyranosyl or C2-C6 open-chain structure.

In this process, acylated (C2 to C3) carbohydrate, preferably, mono to tri-saccharide) is reacted with free fatty acid (C6 to C22) and the reaction temperature ranges from 60 to 95 degree C, preferably from 80 to 90 degree C, in the presence of an acid catalyst, under reduced pressure within the range of 4 to 20 Torr, preferably from 5 to 10 Torr and, without adding any solvent. Next, the reaction mixture is decolorized and separated into unreacted fatty acid layer and a carbohydrate fatty-ester layer.

The unreacted C2-C3 acylated carbohydrate is then precipitated out by cooling the carbohydrate fatty-ester layer to 10 to 0 degree C. If desirable, the free hydroxyl groups are liberated by partial hydrolysis in the presence of an acid or base catalyst. The unreacted free fatty acids and carbohydrate esters of low molecular-




weight carboxylic acids, which are removed during purification, are then reused. The low molecular weight carboxylic acid, which is a by-product of the trans-acidolysis, is trapped and recycled.

- 5 In accordance with the preferred embodiment of the present invention, the process is carried out at low temperature, within the range of 60 to 95 degree C.

The following examples will further illustrate the present
10 invention.

EXAMPLE 1: Preparation of acetylated glucose laurate and the partially deacetylated glucose laurate

0.03 mol (6.00g) of lauric acid (Fluka Chemika, Switzerland) was
15 added into a three-necked, round-bottom flask equipped with a magnetic stirrer, stopcocks, a vacuum take-off line leading to a liquid nitrogen cold-trap and a vacuum pump. 0.01 mol (3.90g) of Glucose pentaacetate, GPA, (Fluka Chemika, Switzerland) was added and the mixture was heated to 80-100°C in an oil bath with continuous stirring until a homogenous solution
20 was formed. 0.1 % w/w (0.01g) of concentrated sulphuric acid was then added and 10-20 Torr pressure was applied. The reaction was continued for 3 to 6 hours. The reaction was then stopped and taken up with 100ml of n-propanol. The solution was then neutralized with 5.0ml aqueous 1M




NaHCO₃, decolourised with 1.0g activated charcoal and filtered warm. (Ethyl acetate or acetonitrile could also be used in place of n-propanol). N-propanol was then removed by vacuum distillation.

Unreacted lauric acid was extracted with 50 X 4 ml hexane. Hexane
5 was vacuum distilled to recover unreacted lauric acid and kept for reuse. The glucose fatty ester was redissolved in warm n-propanol, cool to 0 °C and filtered to give unreacted GPA as precipitate. The filtrate n-Propanol was then vacuum distilled off to give 70-85% pure acetylated glucose laurate. Repeated fatty acid extraction and GPA precipitation gave 90-95% pure
10 product (90.2% yield).

Partial deacetylation of acetylated glucose laurate

Acetylated glucose laurate obtained above was divided into four equal portions. Each portion was refluxed in 150ml 1MHCl (aqueous) solution for various lengths of time, to give partially deacetylated glucose fatty esters of
15 various HLB values, depending on the length of reflux. The solvent was distilled off under reduced pressure.

This Sheet Has Been Replaced



Example 2: Preparation of acetylated sucrose oleate and the partially deacetylated sucrose oleate

0.06mol (16.93g) of oleic acid (Fluka Chemika, Switzerland) was
5 weighed into a three-necked, round-bottom flask equipped with a magnetic
stirrer, stopcocks, a vacuum take-off line leading to a liquid nitrogen cold-
trap and a vacuum pump. 0.01mol (6.79g) of sucrose octaacetate, SOA,
(Fluka Chemika, Switzerland) was added and the mixture was heated to 80-
100°C in an oil bath with continuous stirring until a homogenous solution
10 was formed. 0.1 % w/w (0.024g) of concentrated sulphuric acid (or 0.5g
alkyl sulfonic acid polysiloxane) was then added and 10-20 Torr pressure
was applied. The reaction was continued for 3 to 6 hours. The product was
taken up in 250ml of ethyl acetate, neutralized with 5.0ml aqueous 1M
NaHCO₃, decolourised with 2.0g activated charcoal, and filtered. Ethyl
15 acetate was then removed by vacuum distillation.

Unreacted oleic acid was extracted with 50x4 ml hexane and kept for
reuse. The sucrose fatty ester was redissolved in warm n-propanol, cool to 0
°C and filtered to give unreacted GPA as precipitate. The filtrate n-Propanol
was then distilled off to give 70-85% pure acetylated sucrose oleate.
20 Repeated hexane extraction and SOA precipitation gave 90-95% pure
product (90.2% yield).

This Sheet Has Been Replaced




Partial deacetylation of acetylated sucrose oleate

Acetylated sucrose oleate obtained above was divided into four equal portions. 150 ml aqueous ammonia solution was added to of the portions and
5 agitated for various times, to give sucrose fatty esters of various HLB values, depending on the length of treatment with the ammonia solution. The solvent was distilled off under reduced pressure.

While the invention has been described with respect to
10 preferred embodiment, it will be clear to those skilled in the art that modifications and improvements may be made to the invention without departing from the spirit and scope of the invention. Therefore, the invention is not to be limited by the specific illustrative embodiment, but only by the scope of the appended claims.

15

This Sheet Has Been Replaced





CLAIMS

1. A solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters comprising the steps of:

(a) reacting acylated carbohydrate with free fatty acid in the presence of an acid catalyst, under reduced pressure;

(b) decolorizing and separating the reaction mixture obtained in step (a) into unreacted fatty acid layer and a carbohydrate fatty-ester layer;

(c) precipitating out the unreacted acylated carbohydrate;

(d) liberating the free hydroxyl groups by partial hydrolysis in the presence of an acid or base catalyst;

(e) removing the unreacted free fatty acids and carbohydrate esters of low molecular-weight carboxylic acids during purification, and recycling the removed unreacted free fatty acids and carbohydrate esters to the starting reactant mixture

2. A solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters as set forth in Claim 1, wherein in step (a), no solvent is added thereto.

3. A solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters as set forth in Claim 1, wherein the unreacted acylated carbohydrate is

This Sheet Has Been Replaced





precipitated out by cooling the carbohydrate fatty-ester layer to 10 to 0 degree C.

- 5 4. A solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters as set forth in Claim 1, wherein step (a) is processed at a temperature ranging from 60 to 95 degree C.

10

This Sheet Has Been Replaced



ABSTRACT OF THE DISCLOSURE

A SOLVENT-FREE TRANS-ACIDOLYSIS PROCESS FOR
PREPARING EDIBLE SURFACE-ACTIVE CARBOHYDRATE
FATTY-ACID ESTERS

5

The present invention relates to a low temperature, solvent-free trans-acidolysis process for preparing edible surface-active carbohydrate fatty-acid esters comprising the steps of:

10

(a) reacting acylated carbohydrate with free fatty acid in the presence of an acid catalyst, under reduced pressure;

(b) decolorizing and separating the reaction mixture obtained in step (a) into unreacted fatty acid layer and a carbohydrate fatty-ester layer;

15

(c) precipitating out the unreacted acylated carbohydrate;

(d) liberating the hydroxyl groups by partial hydrolysis in the presence of an acid or base catalyst;

20

(e) removing the unreacted free fatty acids and carbohydrate esters of low molecular-weight carboxylic acids during purification, and recycling the removed unreacted free fatty acids and carbohydrate esters to the starting reactant mixture

This Sheet Has Been Replaced

